



## Monitor Black Layer to Guide Corn Harvest

### KEY POINTS

- Physiological maturity in corn is determined by kernels reaching black layer.
- Timing of corn maturation can be greatly affected by environmental issues.
- Grain moisture content should be monitored once black layer is achieved.
- Harvesting should take place at a moisture content that balances harvest losses and grain drying costs.

Corn kernels achieve black layer or physiological maturity when a black film develops at the tip of a kernel (Figure 1). The black film or layer seals the kernel from further development or increase in test weight. The kernel moisture content at black layer formation usually ranges from 25 to 40 percent, but averages around 30 percent.

Physiological maturity is greatly influenced by an individual product's bred-in relative maturity (RM). Short season products mature earlier than fuller season products. This genetic characteristic allows for the selection and planting of different corn products to spread out (over time) the maturation of a farming operation's corn crop. This helps with harvest scheduling as all corn acres are not maturing at the same time.

Environmental stress can affect the timing of maturation. Severe drought can cause plants to die prematurely and kernels to form a premature black layer. Cool weather during the growing season can delay the normal maturation for an individual product because the growing degree units (GDUs) required to achieve maturity accumulate slower.

Grain moisture content should start to be monitored soon after physiological maturity (black layer). A harvest moisture content level that balances harvest losses and grain drying costs allows maximum economic return potential. A plan to take advantage of a rapid drydown and allow everything to field dry could be costly. Harvesting at lower moistures can increase mechanical losses due to ear drop, stalk lodging, and kernel shattering. Consider beginning harvest when corn grain moisture content is a little above 25% so that harvesting can be finished before corn dries completely in the field.<sup>1</sup>

With high temperatures, it is extremely easy to underestimate the rate that grain dries. Grain that matures in late July can have an average daily drydown rate of approximately 0.8 percentage point per day compared to 0.4 percentage point per day for grain nearing maturity later in the season. Regardless of the environment, it is normal to see later RM corn products reach harvestable moisture levels (around 25%) later than earlier RM products. Product drydown rates are influenced by many product characteristics, such as tightness of husk leaves and pericarp thickness of the seed.<sup>2</sup>

In addition to harvesting at an optimum grain moisture content, achieving proper combine settings can help increase combine efficiency, maximize grain quality, and minimize field losses. Always follow the manufacturer's equipment setting recommendations.




Figure 1. Corn kernel displaying black layer.

Sources: <sup>1</sup>Nielsen, R.L. 2013. Grain fill stages in corn. Corny News Network Articles. Purdue University. <https://www.agry.purdue.edu/ext/corn/news/timeless/GrainFill.html>. <sup>2</sup>Nielsen, R.L. 2013. Field drydown of mature corn grain. Corny News Network Articles. Purdue University. <https://www.agry.purdue.edu/ext/corn/news/timeless/GrainDying.html>. Web sources verified 6/14/18.

Performance may vary, from location to location and from year to year, as local growing, soil and weather conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and should consider the impacts of these conditions on the grower's fields. ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. ©2018 Monsanto Company. All Rights Reserved. 180718135500 061418JMG